

AD-A171 937

PORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION unclass			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited. (2)		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 86-0628		
6a. NAME OF PERFORMING ORGANIZATION Univ of Calif at Davis-Livermore		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR		
6c. ADDRESS (City, State and ZIP Code) Applied Science Dept Livermore CA 94550			7b. ADDRESS (City, State and ZIP Code) same as 6c		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NE	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620-81-C-0050		
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2306 ARPA	TASK NO. B2
11. TITLE (Include Security Classification) FUNDAMENTAL STUDY OF DENSE-FLUID DETONATION			WORK NO.		
12. PERSONAL AUTHOR(S) Professor W.G. HOOVER					
13a. TYPE OF REPORT ANNUAL		13b. TIME COVERED FROM 01Apr83 to 31Mar84		14. DATE OF REPORT (Yr., Mo., Day)	
				15. PAGE COUNT 26	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>OTIC FILE COPY</p> <p>We have determined the atomistic pair distribution function for planar crystals undergoing both uniaxial and hydrostatic compression. We have compared these distributions to the predictions of the Pastine-Kamlet-Jacobs model. We have simulated bimolecular collisions of triatomic and hexatomic planar molecules (with hexanitrobenzene in mind). We analyze the normal-mode vibrations before and after collision and observe the transfer of energy among the translational, rotational, and vibrational modes. During the remainder of this contract we will combine the energy transfer and deformation techniques to study a model of solid hexanitrobenzene.</p> <p>are compared is determined</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL Lt Col Harry V. Winsor			22b. TELEPHONE NUMBER (Include Area Code) 202-767-4931		22c. OFFICE SYMBOL AFOSR/NE

*Covers*

*1 Apr 83-31 Mar 84*

"The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either expressed or implied of the Defense Advanced Research Projects Agency or the U.S. Government."

- (A) ARPA Order 4077
- (B) Program Code 1D10
- (C) Name of Contractor - University of California
- (D) Effective Date of Contract - 1 April 1981
- (E) Contract Expiration Date - 30 September 1985
- (F) Amount of Contract Dollars - \$151,842
- (G) Contract Number - F49620-81-C-0050, P00004
- (H) Principal Investigator - William G. Hoover, (415) 422-9787
- (I) Program Manager-AFOSR/NE, Director of Electronic and Material Sciences  
Building 410, Bolling AFB, DC 20332, (202) 767-4931
- (J) Short Title of Work - Fundamental Detonation Studies

DTIC  
ELECT  
S SEP 15 198  
A

Sponsored by  
Advanced Research Projects Agency (DOD)  
ARPA Order No. 4077  
Monitored by AFOSR Under Contract #F49620-81-C-0050  
Air Force Office of Scientific Research Annual Report - 30 March 1984

Contract F49620-81-C-0500  
81-00645/AO 4077 Effective 1 April 81

Approved for public release;  
distribution unlimited.

"FUNDAMENTAL STUDY OF DENSE-FLUID DETONATION"

Professor W. G. Hoover  
Principal Investigator

Department of Applied Science  
University of California at Davis-Livermore  
Livermore, California 94550  
(415) 422-9787

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DTIC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12.  
Distribution is unlimited.  
W. G. J. KEEPER  
Technical Information Division

## SUMMARY

### (1) TECHNICAL PROBLEM

We are investigating atomistic mathematical models of energy transferred in condensed media. We wish to obtain a fundamental understanding of the interaction of compressive waves with intermolecular and intramolecular energies, leading to the initiation of detonation.

### (2) GENERAL METHODOLOGY

We analyze quasistatic compression by comparing equilibrium statistical-mechanical calculations with the results of equilibrium Newtonian molecular dynamics simulations. We are developing and solving nonequilibrium, equations of motion to describe the rapid compression of microscopic polyatomic systems. We emphasize the flow of energy among the molecular degrees of freedom excited by the compression process.

### (3) TECHNICAL RESULTS

We have determined the atomistic pair distribution function for planar crystals undergoing both uniaxial and hydrostatic compression. We have compared these distributions to the predictions of the Pastine-Kamlet-Jacobs model.

We have simulated bimolecular collisions of triatomic and hexatomic planar molecules (with hexanitrobenzene in mind). We analyze the normal-mode vibrations before and after collision and observe the transfer of energy among the translational, rotational, and vibrational modes.

### (4) FURTHER RESEARCH IMPLICATIONS

The simulations of rapid deformation will be extended to larger molecules, in three dimensions, and will be made consistent with macroscopic thermodynamics. These extensions require equations of motion incorporating a nonequilibrium inhomogeneous deformation satisfying the adiabatic first law  $\dot{E} \equiv -VP:\nabla u$ .

## I. INTRODUCTION

Shock and detonation waves involve atomic-scale processes which are imperfectly understood and hard to measure. The systems contain too many atoms for an accurate quantum-mechanical treatment. Classical calculations can deal with systems of hundreds or thousands of interacting atoms, over hundreds or thousands of vibrational periods.

To evaluate the utility of the microscopic approach we have embarked on a program to develop and apply modern computational methods to the nonequilibrium dynamical processes responsible for explosive initiation. A detailed microscopic view is necessitated by the violent nonequilibrium nature of shock compression. Simulations show that the intrinsic width of a shock front is only a few atomic diameters.<sup>1</sup> Thus the usual concept of equilibrium temperature, with Arrhenius kinetics, is unlikely to provide an understanding of the initiation of chemical reactions.<sup>2</sup> To achieve understanding of initiation detailed studies of rapid nonequilibrium intermolecular and intramolecular energy transfer must be carried out and analyzed.

Over the past several years there has been considerable progress in treating the deformation of simple monatomic fluids and solids. The techniques developed have been tested by intercomparison with experimental data and with less-efficient older simulation techniques.<sup>3</sup> The irreversible heating associated with rapid viscons and plastic deformation has been avoided by developing isothermal equations of motion.<sup>4</sup> These equations constrain the second moment of the velocity distribution,  $\langle(v-\langle v \rangle)^2\rangle$ , to a fixed value. Analogous developments for treating diffusion and conduction stem from this same foundation.<sup>5</sup>

These new methods have not been applied to flexible polyatomic molecules. In order to carry out this generalization we have analyzed both quasistatic and rapid deformations of monatomic crystals (Section II), the rotational and colli-



sional transfer of polyatomic molecular energies (Section III), and the thermodynamic consistency of rapid polyatomic deformation (Section IV). During the remainder of this contract we will combine the energy transfer and deformation techniques to study a model of solid hexanitrobenzene.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A1	

## II. DEFORMATION OF MONATOMIC SOLIDS

In a series of studies we have characterized the compression of simple fluids and solids. We recently carried out a detailed study of equilibrium pair distribution functions in deformed two-dimensional crystals.<sup>6</sup> Both Hooke's-Law and Lennard-Jones forces were used. The pair distributions, obtained by integrating the quasiharmonic canonical probability density, differ from approximate theoretical treatments in two ways. First, contributions from all normal-mode vibrations are included. Second, reaction-rate contributions from all diatomic orientations are included. For these reasons the reaction rates follow neither the simple Gruneisen density dependence nor Arrhenius kinetics. Even under quasistatic conditions, uniaxial compression generally yields significantly higher collision (or reaction) rates than does the corresponding hydrostatic compression.

Earlier deformation studies, under both shockwave and homogeneous conditions, have demonstrated the utility of incorporating the macroscopic velocity gradient  $\nabla u$  directly into microscopic equations of motions. The shockwave studies emphasized the nonequilibrium velocity distribution in the shock-front. Shockwave structure, bulk and shear viscosities, and solid-phase high-strain-rate yield strengths<sup>7</sup> have all been determined in this way. Adiabatic deformation can be modeled by adding an extra contribution to the atomic velocities,

$\dot{q} = q \cdot \nabla u$ , and a corresponding force  $\dot{p} = -\nabla u \cdot p$ . The two modifications lead to the thermodynamic identity

$$\dot{E} = -VP : \nabla u,$$

where  $E$  is internal energy and  $P$  is the pressure tensor. This adiabatic deformation can be made isothermal instead by including an additional collective force  $-\beta p$ , with  $\beta$  chosen to keep  $\langle p^2/m \rangle$  constant.

### III. ENERGY TRANSFER IN POLYATOMIC MOLECULES

The simplest polyatomic molecule exhibiting energy transfer is a planar triatomic molecule, with three vibrational degrees of freedom. In this case the two degenerate vibrational modes are coupled by Coriolis forces. Thus rotation of the molecule, even very slowly, leads to a geometric coupling of two vibrational amplitudes.

Regular hexatomic planar molecules have nine vibrational degrees of freedom, with four pairs of degenerate modes in addition to the symmetric breathing mode. At reasonable rotational velocities the coupling among the four pairs is negligible. Even at vanishing velocities Coriolis coupling again causes the cycling back and forth between degenerate mode amplitudes.

We have simulated bimolecular collisions of both triangular and hexagonal planar molecules, have analyzed their normal-mode vibrations throughout the collision process, and have observed energy transfer among all the translational, rotational, and vibrational degrees of freedom. Presently we are characterizing the energy distribution during a homogeneous compression.

#### IV. THERMODYNAMICS OF RAPID DEFORMATION

The virial theorem relates the pressure tensor to interatomic forces. In the familiar monatomic form<sup>8</sup> each Newton's equation of motion is multiplied by the corresponding particle coordinate, summed and averaged, to give

$$\sum \sum r_{ij} F_{ij} - VP = \sum \frac{d}{dt} (m \mathbf{r} \dot{\mathbf{r}})_i - (m \dot{\mathbf{r}} \mathbf{r})_i.$$

which is equivalent to

$$PV = NkTI + \sum r_{ij} F_{ij}$$

where N is the number of atoms in the volume pairs. For polyatomic molecules it is convenient to multiply each atom's equation of motion by the center of mass coordinate for its molecule. The force terms in each molecule cancel, with the result:

$$PV = NkTI + \sum R_{ij} F_{ij}$$

where N is now the number of molecules and  $F_{ij}$  is the vector sum of forces of all atoms in molecule I due to atoms in molecule J.

Now consider two kinds of adiabatic deformation. In "atomic" deformation each atom undergoes a displacement proportional to its location

$$\dot{\mathbf{r}} = \mathbf{r} \cdot \nabla u$$

and a corresponding force

$$\dot{\mathbf{p}} = -\mathbf{p} \cdot \nabla u.$$

The adiabatic first-law identity,  $\dot{\mathbf{E}} = -VP \cdot \nabla u$ , follows. In "molecular" deformation each atom undergoes a displacement proportional to the center-of-mass coordinate R:

$$\dot{\mathbf{r}} = \mathbf{R} \cdot \nabla u$$

as well as a corresponding force

$$\dot{\mathbf{p}} = -P \left( \frac{\mathbf{r}}{M} \right) \cdot \nabla u.$$

The identity  $\dot{\mathbf{E}} = -PV : \nabla u$  again follows, provided that P is the "molecular" pressure.



It is presently unknown which of these formulations best describes adiabatic polyatomic deformation. We will apply both methods to rapid deformation of hexagonal molecules in our energy transfer simulations in the coming year.

#### REFERENCES:

- (1) B. L. Holian, W. G. Hoover, B. Moran and G. K. Straub, Phys. Rev. A 22, 2798, 1980.
- (2) V. Yu. Klimenko, A. N. Dremin, "The Effect of the Shockwave Front on the Origin of Reaction," Seventh ICOGER, Gottingen--Am. Inst. Aero. and Astro., Inc., 1979.
- (3) W. T. Ashurst, Dense Fluid Shear Viscosity & Thermal Conductivity via Nonequilibrium Molecular Dynamics, Thesis, UC Davis, 1974.
- (4) W. G. Hoover, Atomistic Nonequilibrium Computer Simulations, Physica 118A, 111-122, 1983.
- (5) D. J. Evans, "Computer Experiment for Non-Linear...", J. Chem. Phys., (to be published).
- (6) A. Combs and W. G. Hoover, in preparation.
- (7) W. G. Hoover, A. J. C. Ladd, B. Moran, "High-Strain-Rate Plastic Flow Studied via Nonequilibrium Molecular Dynamics," Phys. Rev. Letts. 48:26, 1982.
- (8) Hirschfelder, Curtiss, and Bird, Molecular Theory of Gases and Liquids, (Wiley, NY).